

**ON INVERSE SOURCE PROBLEM FOR SOBOLEV EQUATION
 WITH MITTAG-LEFFLER KERNEL IN L^r SPACE**

BUI DUC NAM, LE XUAN DAI, LE DINH LONG, NGUYEN HOANG TUAN

ABSTRACT. In this paper, we consider a Sobolev equation with the Atangana-Baleanu-Caputo fractional derivative. We give the explicit fomula of the source term. Under the observations of tha data in L^r spaces, we provide a regularized solution using Fourier truncated method. We give the error estimate between the exact solution and the regularized solution. The main tool is of using some embeddings.

1. INTRODUCTION

Let $\Omega \in \mathbb{R}^N (N \geq 1)$ be a bounded domain with sufficiently smooth boundary $\partial\Omega$. In this paper, we are interested to study time fractional diffusion equation with fractional derivative as follows

$$\begin{cases} {}_0^{ABC}D_t^\beta (u(x, t) + m\mathcal{L}u(x, t)) + \mathcal{L}u(x, t) = g(t)f(x), & x \in \Omega, t \in (0, 1), \\ u(x, t) = 0, & x \in \partial\Omega, t \in (0, 1), \\ u(x, 1) = \rho(x), & x \in \Omega. \end{cases} \quad (1.1)$$

Here in the main equation as above, the Atangana - Baleanu fractional derivative ${}_0^{ABC}D_t^\beta u(x, t)$ is defined by

$${}_0^{ABC}D_t^\beta u(x, t) = \frac{\mathcal{M}(\beta)}{1 - \beta} \int_0^t \frac{\partial u(x, s)}{\partial s} E_{\beta, 1} \left(\frac{-\beta(t - s)^\beta}{1 - \beta} \right) ds, \quad (1.2)$$

where the normalization function $\mathcal{M}(\beta)$ can be any function satisfying the conditions $\mathcal{M}(\beta) = 1 - \beta + \frac{\beta}{\Gamma(\beta)}$, here $\mathcal{M}(0) = \mathcal{M}(1) = 1$ (see Definition 2.1 in [1]) and $E_{\beta, 1}$ is the MittagLeffler function. Our main goal in this paper is of finding the source term $f(x)$ from the given data ρ and the measured data at the final time

2000 *Mathematics Subject Classification.* 35K05, 35K99, 47J06, 47H10x .

Key words and phrases. Source problem; Fractional pseudo-parabolic problem; Ill-posed problem; Convergence estimates; Regularization, Mittag-Leffler kernel, Atagana-Baleanu-Caputo derivative.

©2023 Universiteti i Prishtinës, Prishtinë, Kosovë.

Submitted September 14, 2023. Published October 31, 2023.

Communicated by Douglas R. Anderson.

$u(x, 1) = \rho(x)$, $\rho \in L^2(\Omega)$ such that

$$\|g - g_\epsilon\|_{L^r(0,1)} + \|\rho - \rho_\epsilon\|_{L^r(\Omega)} \leq \epsilon. \quad (1.3)$$

One of the narrow branches of fractional analysis is the theory of fractional diffusion equations. Fractional-time diffusion equations are used to model complex phenomena such as long-term memory or spatial interactions, non-local and local dynamics. For details, please refer to documents [2, 3, 5, 6, 7, 8, 18]. One of the modern trends in fractional analysis is the development of fractional operators with non-singular kernels. The study of these fractional derivatives is important to satisfy the need for modeling applications in various fields, such as fluids, mechanics, viscoelasticity, biology, physics and engineering, see in [3, 4, 17, 9, 10, 11, 12, 13, 21, 22, 23, 24]. Several definitions of fractional derivatives have been given based on non-special nuclei such as Atangana-Baleanu-Caputo fractions and derivatives. Regarding the study of (1.1) problem with Atangana-Baleanu derivative, we list some previous results as follows

- Under the case $F = F(x, t)$, from [19], the kernels of the extended Mittag-Leffler type functions are studied in this study using a partial differential equation model with the new universal fractional derivatives. Analysis and consideration are given to an initial boundary value problem for the anomalous diffusion of fractional order. The Mittag-Leffler kernel fractional derivative, also known as the Atangana-Baleanu fractional derivative in time, is interpreted in the Caputo sense. They discovered findings on the existence, uniqueness, and regularity of the solution.
- Under the case $F = g(t)f(x)$, from [14, 16], the problem of determining inverse source problem for fractional diffusion equation containing Atangana-Baleanu-Caputo fractional derivative. We first establish an explicit formula of the source term from the average data of the function in the time variable. We then show that the inverse source problem is ill-posed in the meaning of Hadamard i.e., the source function is not stable according to the given data. To overcome this instability, we propose a regularized method as in the Fractional Landweber method. We also obtain the upper bounds and find the convergence rate between the regularized solution and sought source function. Estimates are also derived in two cases on selection rules, a priori parameter, and a posterior parameter. Numerical examples are given which illustrate the usefulness of our method.
- Under the case $F = F(x, t, u(x, t))$, in the paper [15], they investigated a nonlinear time fraction Volterra equation with a Mittag-Leffler multiplier in Hilbert space. By applying the properties of the Mittag-Leffler function and the eigenvalue expansion, the existence of a light solution to our problem has been proved. The main tool to prove our results is the use of some Sobolev embeddings and some fixed point theorems.

As we know, the inverse issue for diffusion equation with Atangana-Baleanu fractional derivative where the observed data is in the $L^p(\Omega)$ space with $p \neq 2$ is solved for the first time in this study. One significant challenge is that because the data is not in $L^2(\Omega)$, we cannot utilize Parseval equality directly. We get around these issues by leveraging embedding between $L^p(\Omega)$ and $\mathcal{H}^s(\Omega)$ scale-spaces. We have the regularized solution through the Fourier series truncation method with the observed data $(\varphi_\epsilon, g_\epsilon) \in L^r(0, T) \times L^r(\Omega)$. After that, the error established between

the regularized solution and the exact solution in the Theorem (3.1), by the main analytical technique is to use some embeddings and some evaluations using Hlder inequality.

The structure of our paper is as follows. The existence of mild solution u to (1.1) in Section 1. In the Section 2, we have some preliminaries. The main results in Section 3 is Theorem 3.1, our main tool is Sobolev embeddings.

2. PRELIMINARIES

Let us recall that the spectral problem

$$\begin{cases} -\mathcal{L}\varphi_n(x) = \xi_n\varphi_n(x), & \text{in } \Omega, \\ \varphi_n(x) = 0, & \text{on } \partial\Omega, \end{cases} \quad (2.1)$$

admits a family of eigenvalues

$$0 < \xi_1 \leq \xi_2 \leq \xi_3 \leq \dots \leq \xi_n \leq \dots \nearrow \infty.$$

For all $r \geq 0$, the operator \mathcal{L}^r (here $\mathcal{L} = -\Delta$) also possesses the following representation:

$$\begin{aligned} \mathcal{L}^r h &= \sum_{n=1}^{\infty} \left(\int_{\Omega} h(x)\varphi_n(x)dx \right) \xi_n^r \varphi_n, \\ h \in \mathcal{H}^r(\Omega) &= \left\{ h \in L^2(\Omega) : \sum_{n=1}^{\infty} \left(\int_{\Omega} h(x)\varphi_n(x)dx \right)^2 \xi_n^{2r} < \infty \right\}. \end{aligned} \quad (2.2)$$

Consider on \mathcal{H}^r the norm

$$\|h\|_{\mathcal{H}^r(\Omega)} = \left(\sum_{n=1}^{\infty} \left(\int_{\Omega} h(x)\varphi_n(x)dx \right)^2 \xi_n^{2r} \right)^{\frac{1}{2}}, \quad h \in \mathcal{H}(\mathcal{A}^r).$$

Lemma 2.1. ([6]) *Let $0 < \beta < 1$, then there exist $0 < \mathcal{B}_1, \mathcal{B}_2, \mathcal{B}_3$ such that*

$$\frac{\mathcal{B}_1}{1+y} \leq E_{\beta,1}(-y) \leq \frac{\mathcal{B}_2}{1+y}, \quad E_{\beta,\alpha}(-y) \leq \frac{\mathcal{B}_3}{1+y}, \quad \text{for all } y \geq 0, \alpha \in \mathbb{R}. \quad (2.3)$$

Lemma 2.2. *For $\xi > 0, \beta > 0, m \in \mathbb{N}^*$, we have*

$$\begin{aligned} \frac{d^m}{dt^m} E_{\beta,1}(-\xi t^\beta) &= -\xi t^{\beta-m} E_{\beta,\beta-m+1}(-\xi t^\beta), \\ \frac{d}{dt} (t E_{\beta,2}(-\xi t^\beta)) &= E_{\beta,1}(-\xi t^\beta), \\ \frac{d}{dt} (t^{\beta-1} E_{\beta,\beta}(-\xi t^\beta)) &= -t^{\beta-2} E_{\beta,\beta-1}(-\xi t^\beta). \end{aligned} \quad (2.4)$$

Lemma 2.3. [6] *For $t > 0$, and $\xi > 0$, and $0 < \beta < 1$, then one has*

$$\partial_t^\beta E_{\beta,1}(-\xi t^\beta) = -\xi E_{\beta,1}(-\xi t^\beta),$$

Lemma 2.4. *For $\beta \in (0, 1)$, putting $\mathcal{A}_{3,n}(m, \beta) = \frac{\mathcal{M}(\beta)}{\xi_n(\mathcal{M}(\beta) + \xi_n \sigma_n(1 - \beta))}$, it gives*

$$\mathcal{A}_{3,n}(m, \beta) \geq \frac{1}{\xi_n} \frac{m\mathcal{M}(\beta)}{\left(\frac{\mathcal{M}(\beta)}{\xi_1 \sigma_1} + (1 - \beta) \right)}. \quad (2.5)$$

Proof. First of all, we notice that $\xi_n \sigma_n = \frac{\xi_n}{1+m\xi_n} \geq \frac{1}{\xi_1^{-1}+m} = \frac{1}{\xi_1^{-1}+m}$, and we have

$$\begin{aligned} \mathcal{A}_{3,n}(m, \beta) &= \frac{\mathcal{M}(\beta)}{\xi_n(\mathcal{M}(\beta) + \xi_n \sigma_n(1-\beta))} \geq \frac{\mathcal{M}(\beta)}{\frac{\xi_n^2}{1+m\xi_n} \left(\frac{\mathcal{M}(\beta)}{\xi_n \sigma_n} + (1-\beta) \right)} \\ &\geq \frac{m\mathcal{M}(\beta)}{\xi_n \left(\frac{\mathcal{M}(\beta)}{\xi_n \sigma_n} + (1-\beta) \right)} \geq \frac{1}{\xi_n} \frac{m\mathcal{M}(\beta)}{\left(\frac{\mathcal{M}(\beta)}{\xi_1 \sigma_1} + (1-\beta) \right)}. \end{aligned} \quad (2.6)$$

□

Lemma 2.5. For $M > 0$, by Lemmas 2.1 and 2.2, we have

$$\begin{aligned} \int_0^M \left| t^{\alpha-1} E_{\alpha,\alpha}(-\lambda_n t^\alpha) \right| dt &= \int_0^M t^{\alpha-1} E_{\alpha,\alpha}(-\lambda_n t^\alpha) dt \\ &= -\frac{1}{\lambda_n} \int_0^n \frac{d}{dt} E_{\alpha,1}(-\lambda_n t^\alpha) dt = \frac{1}{\lambda_n} (1 - E_{\alpha,1}(-\lambda_n M^\alpha)). \end{aligned} \quad (2.7)$$

Lemma 2.6. Let $\beta \in (0, 1)$, we have estimate

$$\begin{aligned} &\frac{1}{\xi_n} \frac{m\mathcal{M}(\beta)}{\left(\frac{\mathcal{M}(\beta)}{\xi_1 \sigma_1} + (1-\beta) \right)} \left[1 - E_{\beta,1} \left(-\frac{\beta(\xi_1^{-1} + m)^{-1}}{\mathcal{M}(\beta) + (\xi_1^{-1} + m)^{-1}(1-\beta)} \right) \right] \\ &\leq \frac{\sigma_n \beta \mathcal{M}(\beta)}{(\mathcal{M}(\beta) + \sigma_n \xi_n(1-\beta))^2} \int_0^1 E_{\beta,\beta} \left(-\frac{\beta \xi_n \sigma_n (1-s)^\beta}{\mathcal{M}(\beta) + \sigma_n \xi_n(1-\beta)} \right) (1-s)^{\beta-1} ds \leq \frac{1}{\xi_n}. \end{aligned} \quad (2.8)$$

Proof. For $E_{\beta,\beta}(-z) \geq 0$ for $0 < \beta < 1$ and $z \geq 0$, and using the Lemmas 2.4 and 2.6, we obtain

$$\begin{aligned} a) &\frac{\sigma_n \beta \mathcal{M}(\beta)}{(\mathcal{M}(\beta) + \sigma_n \xi_n(1-\beta))^2} \int_0^1 E_{\beta,\beta} \left(-\frac{\beta \xi_n \sigma_n (1-s)^\beta}{\mathcal{M}(\beta) + \sigma_n \xi_n(1-\beta)} \right) (1-s)^{\beta-1} ds \\ &\geq \frac{1}{\xi_n} \frac{m\mathcal{M}(\beta)}{\left(\frac{\mathcal{M}(\beta)}{\xi_1 \sigma_1} + (1-\beta) \right)} \left[1 - E_{\beta,1} \left(-\frac{\beta(\xi_1^{-1} + m)^{-1}}{\mathcal{M}(\beta) + (\xi_1^{-1} + m)^{-1}(1-\beta)} \right) \right]. \end{aligned} \quad (2.9)$$

$$\begin{aligned} b) &\frac{\sigma_n \beta \mathcal{M}(\beta)}{(\mathcal{M}(\beta) + \sigma_n \xi_n(1-\beta))^2} \int_0^1 E_{\beta,\beta} \left(-\frac{\beta \xi_n \sigma_n (1-s)^\beta}{\mathcal{M}(\beta) + \sigma_n \xi_n(1-\beta)} \right) (1-s)^{\beta-1} ds \\ &= -\frac{\mathcal{M}(\beta)}{\xi_n(\mathcal{M}(\beta) + \xi_n \sigma_n(1-\beta))} \int_0^1 \frac{d}{ds} \left(E_{\beta,1} \left(-\frac{\beta \xi_n \sigma_n (1-s)^\beta}{\mathcal{M}(\beta) + \sigma_n \xi_n(1-\beta)} \right) \right) d\tau \\ &= \frac{1}{\xi_n} \left(1 - E_{\beta,1} \left(-\frac{\beta \xi_n \sigma_n}{\mathcal{M}(\beta) + \sigma_n \xi_n(1-\beta)} \right) \right) \leq \frac{1}{\xi_n}. \end{aligned} \quad (2.10)$$

□

Lemma 2.7. [17] *The following statement are true:*

$$\left. \begin{aligned} L^r(\Omega) &\hookrightarrow \mathcal{H}^s(\Omega), & \text{if } & -\frac{N}{4} < s \leq 0, & r &\geq \frac{2N}{N-4s}, \\ \mathcal{H}^s(\Omega) &\hookrightarrow L^r(\Omega), & \text{if } & 0 \leq s < \frac{N}{4}, & r &\leq \frac{2N}{N-4s}. \end{aligned} \right\} \quad (2.11)$$

3. THE INVERSE SOURCE PROBLEM (1.1)

Let us first to review the initial value problem as follows

$$\begin{cases} {}_0^{ABC}D_t^\beta(u(x,t) + m\mathcal{L}u(x,t)) + \mathcal{L}u(x,t) = F(x,t), & \text{in } \Omega \times (0, 1], \\ u(x,t) = 0, & \text{on } \partial\Omega \times (0, 1], \\ u(x,0) = u_0(x), & \text{in } \Omega, \end{cases} \quad (3.1)$$

where u_0 and F are given functions. Let $u(x,t) = \sum_{j=1}^{\infty} u_n(t)\varphi_n(x)$ be the Fourier series in $L^2(\Omega)$ with $u_n(t) = \int_{\Omega} u(x,t)\varphi_n(x)dx$, then we have the fractional integro-differential equation involving the Atangana-Baleanu fractional derivative in the form

$${}_0^{ABC}D_t^\beta(1 + m\xi_n)u_n(t) + \xi_n u_n(t) = F_n(t), \quad (3.2)$$

in [20], the solution (3.1) can be represented as by Fourier series

$$u(x,t) = \sum_{n=1}^{\infty} \left(\int_{\Omega} u(x,t)\varphi_n(x)dx \right) \varphi_n(x)$$

and then given by

$$\begin{aligned} u_n(t) &= \frac{\mathcal{M}(\beta)}{\mathcal{M}(\beta) + \frac{\xi_n}{1+m\xi_n}(1-\beta)} E_{\beta,1} \left(\frac{-\beta \frac{\xi_n}{1+m\xi_n} t^\beta}{\mathcal{M}(\beta) + \frac{\xi_n}{1+m\xi_n}(1-\beta)} \right) u_{0,n} \\ &+ \left(\frac{1}{1+m\xi_n} \right) \frac{1-\beta}{\mathcal{M}(\beta) + \frac{\xi_n}{1+m\xi_n}(1-\beta)} F_n(t) \\ &+ \left(\frac{1}{1+m\xi_n} \right) \frac{\beta \mathcal{M}(\beta)}{\left(\mathcal{M}(\beta) + \frac{\xi_n}{1+m\xi_n}(1-\beta) \right)^2} \\ &\times \int_0^t E_{\beta,\beta} \left(\frac{-\beta \frac{\xi_n}{1+m\xi_n} (t-s)^\beta}{\mathcal{M}(\beta) + \frac{\xi_n}{1+m\xi_n}(1-\beta)} \right) (t-s)^{\beta-1} F_n(\tau) d\tau. \end{aligned} \quad (3.3)$$

Let us denote $\sigma_n = (1 + m\xi_n)^{-1}$, this implies that

$$\begin{aligned}
u(x, t) &= \sum_{n=1}^{\infty} \frac{\mathcal{M}(\beta)}{\mathcal{M}(\beta) + \xi_n \sigma_n (1 - \beta)} E_{\beta,1} \left(\frac{-\beta \xi_n \sigma_n t^\beta}{\mathcal{M}(\beta) + \xi_n \sigma_n (1 - \beta)} \right) \langle u(0), \varphi_n \rangle \varphi_n(x), \\
&+ \sum_{n=1}^{\infty} \frac{\sigma_n (1 - \beta)}{\mathcal{M}(\beta) + \xi_n \sigma_n (1 - \beta)} \langle F(\cdot, t), \varphi_n \rangle \varphi_n(x) \\
&+ \sum_{n=1}^{\infty} \frac{\sigma_n \beta \mathcal{M}(\beta)}{(\mathcal{M}(\beta) + \xi_n \sigma_n (1 - \beta))^2} \\
&\quad \times \int_0^t E_{\beta,\beta} \left(\frac{-\beta \xi_n \sigma_n (t - \tau)^\beta}{\mathcal{M}(\beta) + \xi_n \sigma_n (1 - \beta)} \right) (t - s)^{\beta-1} \langle F(\cdot, s), \varphi_n \rangle ds \varphi_n(x). \quad (3.4)
\end{aligned}$$

Let us now return the problem of identifying the source term. Let $t = 1, u(x, 0) = 0, F(x, t) = g(t)f(x)$, and $F_n(1) = 0$, we get

$$\begin{aligned}
\int_{\Omega} \rho(x) \varphi_n(x) dx &= \int_{\Omega} f(x) \varphi_n(x) dx \frac{\sigma_n \beta \mathcal{M}(\beta)}{(\mathcal{M}(\beta) + \xi_n \sigma_n (1 - \beta))^2} \\
&\quad \times \int_0^1 E_{\beta,\beta} \left(\frac{-\beta \xi_n \sigma_n (1 - s)^\beta}{\mathcal{M}(\beta) + \xi_n \sigma_n (1 - \beta)} \right) (1 - s)^{\beta-1} g(s) ds. \quad (3.5)
\end{aligned}$$

To make the formula even more compact, we put

$$\begin{aligned}
\mathcal{A}_{1,n}(m, \beta) &= \frac{\sigma_n \beta \mathcal{M}(\beta)}{(\mathcal{M}(\beta) + \xi_n \sigma_n (1 - \beta))^2}, \\
\mathcal{A}_{2,n}(m, \beta) &= \frac{\beta \xi_n \sigma_n}{\mathcal{M}(\beta) + \xi_n \sigma_n (1 - \beta)}. \quad (3.6)
\end{aligned}$$

From (3.5) and (3.6), we receive

$$\begin{aligned}
\int_{\Omega} \rho(x) \varphi_n(x) dx &= \int_{\Omega} f(x) \varphi_n(x) dx \\
&\quad \times \mathcal{A}_{1,n}(m, \beta) \int_0^1 E_{\beta,\beta} \left(-\mathcal{A}_{2,n}(m, \beta) (1 - s)^\beta \right) (1 - s)^{\beta-1} g(s) ds. \quad (3.7)
\end{aligned}$$

From (3.7), it gives

$$\int_{\Omega} f(x) \varphi_n(x) dx = \frac{\int_{\Omega} \rho(x) \varphi_n(x) dx}{\mathcal{A}_{1,n}(m, \beta) \int_0^1 E_{\beta,\beta} \left(-\mathcal{A}_{2,n}(m, \beta) (1 - s)^\beta \right) (1 - s)^{\beta-1} g(s) ds}. \quad (3.8)$$

Through some basic transformations, we get

$$f(x) = \sum_{n=1}^{\infty} \frac{\int_{\Omega} \rho(x) \varphi_n(x) dx}{\mathcal{A}_{1,n}(m, \beta) \int_0^1 E_{\beta,\beta} \left(-\mathcal{A}_{2,n}(m, \beta) (1 - s)^\beta \right) (1 - s)^{\beta-1} g(s) ds} \varphi_n(x). \quad (3.9)$$

As $n \rightarrow \infty$, i.e., $\left(\mathcal{A}_{1,n}(m, \beta) \int_0^1 E_{\beta, \beta}(-\mathcal{A}_{2,n}(m, \beta)(1-s)^\beta)(1-s)^{\beta-1} g(s) ds\right)^{-1} \rightarrow \infty$, see in the Lemma 2.6. Thus, it can be concluded from formula (3.9) that the small perturbation of $\rho_\epsilon(x)$ will cause a great change of $f(x)$. Thus our problem (1.1) is ill-posed. Next, we will give the conditional stability results of the source term $f(x)$.

Theorem 3.1. *Let us take $(g_\epsilon, \rho_\epsilon) \in L^r(0, 1) \times L^r(\Omega)$ such that $g_\epsilon(t) > G_2 > 0$ for any $0 \leq t \leq 1$ for any $\frac{1}{\beta} < r < 2$ and condition*

$$\|g_\epsilon - g\|_{L^r(0,1)} + \|\rho_\epsilon - \rho\|_{L^r(\Omega)} \leq \epsilon. \quad (3.10)$$

Assume that $f \in \mathcal{H}(\mathcal{A}^{j+k})$ for $k > 0$ and $0 < j < \frac{N}{4}$. With the Fourier truncation method, we have

$$f_\epsilon^{\mathcal{C}_\epsilon}(x) = \sum_{\xi_n \leq \mathcal{C}_\epsilon} \frac{\int_\Omega \rho_\epsilon(x) \varphi_n(x) dx}{\mathcal{A}_{1,n}(m, \beta) \int_0^1 E_{\beta, \beta}(-\mathcal{A}_{2,n}(m, \beta)(1-s)^\beta)(1-s)^{\beta-1} g_\epsilon(s) ds} \varphi_n(x). \quad (3.11)$$

Then we have

$$\begin{aligned} \|f_\epsilon^{\mathcal{C}_\epsilon} - f\|_{L^{\frac{2N}{N-4j}}(\Omega)} &\lesssim |\mathcal{C}_\epsilon|^{-k} \|f\|_{\mathcal{H}^{j+k}(\Omega)} + \mathcal{C}_\epsilon \epsilon |\mathcal{A}_4(\mathcal{B}_3, r, \beta, G_2, m, \xi_1)| \|f\|_{\mathcal{H}^j(\Omega)} \\ &\quad + \mathcal{A}_5(\xi_1, \sigma_1, m, \beta) (\mathcal{C}_\epsilon)^{j+1+\frac{N}{2r}-\frac{N}{4}} \epsilon, \end{aligned} \quad (3.12)$$

whereby \mathcal{C}_ϵ satisfies that

$$\lim_{\epsilon \rightarrow 0} \mathcal{C}_\epsilon \epsilon = \lim_{\epsilon \rightarrow 0} \left((\mathcal{C}_\epsilon)^{j+1+\frac{N}{2r}-\frac{N}{4}} \epsilon \right) = 0, \quad \lim_{\epsilon \rightarrow 0} \mathcal{C}_\epsilon = +\infty. \quad (3.13)$$

Remark. We can take \mathcal{C}_ϵ satisfying (3.13) as follows

$$\mathcal{C}_\epsilon = \epsilon^{\frac{s-1}{j+1+\frac{N}{2r}-\frac{N}{4}}}, \quad 0 < s < 1.$$

Then the error order $\|f_\epsilon^{\mathcal{C}_\epsilon} - f\|_{L^{\frac{2N}{N-4j}}(\Omega)}$ is of order

$$\max \left\{ \epsilon^{\frac{k(j+1+\frac{N}{2r}-\frac{N}{4})}{s-1}}, \epsilon^{\frac{s+j+\frac{N}{2s}-\frac{N}{4}}{j+1+\frac{N}{2s}-\frac{N}{4}}}, \epsilon^s \right\}.$$

Proof. Using the triangle inequality, we have

$$\|f_\epsilon^{\mathcal{C}_\epsilon} - f\|_{\mathcal{H}^j(\Omega)} \leq \underbrace{\|\mathcal{F}_{2,\epsilon} - f\|_{\mathcal{H}^j(\Omega)}} + \underbrace{\|\mathcal{F}_{1,\epsilon} - \mathcal{F}_{2,\epsilon}\|_{\mathcal{H}^j(\Omega)}} + \underbrace{\|\mathcal{F}_{1,\epsilon} - f_\epsilon\|_{\mathcal{H}^j(\Omega)}}, \quad (3.14)$$

where we denote some following functions

$$\mathcal{F}_{1,\epsilon}(x) = \sum_{\xi_n \leq \mathcal{C}_\epsilon} \frac{\int_\Omega \rho(x) \varphi_n(x) dx}{\mathcal{A}_{1,n}(m, \beta) \int_0^1 E_{\beta, \beta}(-\mathcal{A}_{2,n}(m, \beta)(1-s)^\beta)(1-s)^{\beta-1} g_\epsilon(s) ds} \varphi_n(x), \quad (3.15)$$

and

$$\mathcal{F}_{2,\epsilon}(x) = \sum_{\xi_n \leq \mathcal{C}_\epsilon} \frac{\int_{\Omega} \rho(x) \varphi_n(x) dx}{\mathcal{A}_{1,n}(m, \beta) \int_0^1 E_{\beta, \beta}(-\mathcal{A}_{2,n}(m, \beta)(1-s)^\beta) (1-s)^{\beta-1} g(s) ds} \varphi_n(x). \quad (3.16)$$

Let us next consider some terms on the right hand side of (3.14).

Step 1. Estimate of $\|\mathcal{F}_{2,\epsilon} - f\|_{\mathcal{H}^j(\Omega)}$.
Let us recall the function f as follows.

$$f(x) = \sum_{n=1}^{\infty} \frac{\int_{\Omega} \rho(x) \varphi_n(x) dx}{\mathcal{A}_{1,n}(m, \beta) \int_0^1 E_{\beta, \beta}(-\mathcal{A}_{2,n}(m, \beta)(1-s)^\beta) (1-s)^{\beta-1} g(s) ds} \varphi_n(x).$$

This expression together with the fomula (3.16) gives us the claim of the following difference

$$\begin{aligned} & f(x) - \mathcal{F}_{2,\epsilon}(x) \\ &= \sum_{\xi_n \geq \mathcal{C}_\epsilon} \frac{\int_{\Omega} \rho(x) \varphi_n(x) dx}{\mathcal{A}_{1,n}(m, \beta) \int_0^1 E_{\beta, \beta}(-\mathcal{A}_{2,n}(m, \beta)(1-s)^\beta) (1-s)^{\beta-1} g(s) ds} \varphi_n(x) \\ &= \sum_{\xi_n \geq \mathcal{C}_\epsilon} \left(\int_{\Omega} f(x) \varphi_n(x) dx \right) \varphi_n(x). \end{aligned} \quad (3.17)$$

The norm on $\mathcal{H}^j(\Omega)$ of (3.17) is calculated through the Parseval equality as follows

$$\begin{aligned} \|f - \mathcal{F}_{2,\epsilon}\|_{\mathcal{H}^j(\Omega)}^2 &= \sum_{\xi_n \geq \mathcal{C}_\epsilon} \xi_n^{2j} \left(\int_{\Omega} f(x) \varphi_n(x) dx \right)^2 \\ &= \sum_{\xi_n \geq \mathcal{C}_\epsilon} \xi_n^{-2k} \xi_n^{2j+2k} \left(\int_{\Omega} f(x) \varphi_n(x) dx \right)^2. \end{aligned}$$

It is easy to see that $\xi_n^{-2k} \leq |\mathcal{C}_\epsilon|^{-2k}$ if $\xi_n > \mathcal{C}_\epsilon$ and $k > 0$. Hence, we have

$$\begin{aligned} \|f - \mathcal{F}_{2,\epsilon}\|_{\mathcal{H}^j(\Omega)}^2 &\leq |\mathcal{C}_\epsilon|^{-2k} \sum_{\xi_n \geq \mathcal{C}_\epsilon} \xi_n^{2j+2k} \left(\int_{\Omega} f(x) \varphi_n(x) dx \right)^2 \\ &= |\mathcal{C}_\epsilon|^{-2k} \|f\|_{\mathcal{H}^{j+k}(\Omega)}^2, \end{aligned} \quad (3.18)$$

It gives that

$$\|f - \mathcal{F}_{2,\epsilon}\|_{\mathcal{H}^j(\Omega)} \leq |\mathcal{C}_\epsilon|^{-k} \|f\|_{\mathcal{H}^{j+k}(\Omega)}. \quad (3.19)$$

Step 2. Estimate of $\|\mathcal{F}_{1,\epsilon} - \mathcal{F}_{2,\epsilon}\|_{\mathcal{H}^j(\Omega)}$.

Based on two formulas (3.15) and (3.16), we have

$$\begin{aligned} & \mathcal{F}_{1,\epsilon}(x) - \mathcal{F}_{2,\epsilon}(x) \\ &= \sum_{\xi_n \leq \mathcal{C}_\epsilon} \frac{\int_0^1 E_{\beta,\beta}(-\mathcal{A}_{2,n}(m,\beta)(1-s)^\beta)(1-s)^{\beta-1}(g_\epsilon(s) - g(s)) ds}{\int_0^1 E_{\beta,\beta}(-\mathcal{A}_{2,n}(m,\beta)(1-s)^\beta)(1-s)^{\beta-1}g_\epsilon(s) ds} \\ & \quad \times \frac{\int_\Omega \rho(x)\varphi_n(x) dx}{\mathcal{A}_{1,n}(m,\beta) \int_0^1 E_{\beta,\beta}(-\mathcal{A}_{2,n}(m,\beta)(1-s)^\beta)(1-s)^{\beta-1}g(s) ds} \varphi_n(x). \end{aligned} \quad (3.20)$$

We follows from (3.20) that

$$\begin{aligned} & \mathcal{F}_{1,\epsilon}(x) - \mathcal{F}_{2,\epsilon}(x) \\ &= \sum_{\xi_n \leq \mathcal{C}_\epsilon} \frac{\int_0^1 E_{\beta,\beta}(-\mathcal{A}_{2,n}(m,\beta)(1-s)^\beta)(1-s)^{\beta-1}(g_\epsilon(s) - g(s)) ds}{\int_0^1 E_{\beta,\beta}(-\mathcal{A}_{2,n}(m,\beta)(1-s)^\beta)(1-s)^{\beta-1}g_\epsilon(s) ds} \\ & \quad \times \left(\int_\Omega f(x)\varphi_n(x) dx \right) \varphi_n(x). \end{aligned} \quad (3.21)$$

By taking the norm of (3.21) in space $\mathcal{H}^j(\Omega)$ and using Parseval' s equality, we provide that

$$\begin{aligned} & \|\mathcal{F}_{1,\epsilon} - \mathcal{F}_{2,\epsilon}\|_{\mathcal{H}^j(\Omega)}^2 \\ &= \sum_{\xi_n \leq \mathcal{C}_\epsilon} \left[\frac{\int_0^1 E_{\beta,\beta}(-\mathcal{A}_{2,n}(m,\beta)(1-s)^\beta)(1-s)^{\beta-1}(g_\epsilon(s) - g(s)) ds}{\int_0^1 E_{\beta,\beta}(-\mathcal{A}_{2,n}(m,\beta)(1-s)^\beta)(1-s)^{\beta-1}g_\epsilon(s) ds} \right]^2 \\ & \quad \times \xi_n^{2j} \left(\int_\Omega f(x)\varphi_n(x) dx \right)^2. \end{aligned} \quad (3.22)$$

From (3.22), noting that $r > \beta^{-1}$ and $r^* = 1 + \frac{1}{r-1}$, using Hölder inequality and Lemma 3.10, we have

$$\begin{aligned} & \left| \int_0^1 E_{\beta,\beta}(-\mathcal{A}_{2,n}(m,\beta)(1-s)^\beta)(1-s)^{\beta-1}(g_\epsilon(\tau) - g(\tau)) d\tau \right| \\ & \leq \left[\int_0^1 |g_\epsilon(s) - g(s)|^r ds \right]^{\frac{1}{r}} \left[\int_0^1 |E_{\beta,\beta}(-\mathcal{A}_{2,n}(1-s)^\beta)(1-s)^{\beta-1}|^{r^*} ds \right]^{\frac{1}{r^*}} \\ & \leq \|g_\epsilon - g\|_{L^r(0,1)} \left(\mathcal{B}_3^{\frac{r}{r-1}} \frac{r-1}{\beta r - 1} \right)^{\frac{r-1}{r}} = \|g_\epsilon - g\|_{L^r(0,1)} \mathcal{B}_3 \left(\frac{r-1}{\beta r - 1} \right)^{\frac{r-1}{r}}. \end{aligned} \quad (3.23)$$

This implies that

$$\left| \int_0^1 E_{\beta,\beta}(-\mathcal{A}_{2,n}(m,\beta)(1-s)^\beta)(1-s)^{\beta-1}(g_\epsilon(s)-g(s))ds \right| \leq \mathcal{B}_3 \left(\frac{r-1}{\beta r-1} \right)^{\frac{r-1}{r}} \epsilon. \quad (3.24)$$

It is easy to see that

$$\frac{1}{\mathcal{A}_{2,n}(m,\beta)} = \frac{\mathcal{M}(\beta) + \xi_n \sigma_n (1-\beta)}{\beta \xi_n \sigma_n} \geq \frac{\mathcal{M}(\beta)}{\beta \xi_n \sigma_n} \geq \frac{1}{\xi_n} \left(\frac{\mathcal{M}(\beta)}{\beta} \right). \quad (3.25)$$

Next, the function $g_\epsilon \geq G_2$, and using the Lemma 2.6, we have

$$\begin{aligned} & \int_0^1 E_{\beta,\beta}(-\mathcal{A}_{2,n}(m,\beta)(1-s)^\beta)(1-s)^{\beta-1}g_\epsilon(s)ds \\ & \geq G_2 \int_0^1 E_{\beta,\beta}(-\mathcal{A}_{2,n}(1-s)^\beta)(1-s)^{\beta-1}ds \\ & = \frac{G_2}{\xi_n} \frac{\mathcal{M}(\beta)}{\beta} \left[1 - E_{\beta,1} \left(- \frac{\beta(\xi_1^{-1} + m)^{-1}}{\mathcal{M}(\beta) + (\xi_1^{-1} + m)^{-1}(1-\beta)} \right) \right]. \end{aligned} \quad (3.26)$$

From (3.24) and (3.26), we assert that

The right hand side of (3.22)

$$\leq \underbrace{\xi_n \epsilon \mathcal{B}_3 \left(\frac{r-1}{\beta r-1} \right)^{\frac{r-1}{r}} \frac{\beta}{G_2 \mathcal{M}(\beta)} \left[1 - E_{\beta,1} \left(- \frac{\beta(\xi_1^{-1} + m)^{-1}}{\mathcal{M}(\beta) + (\xi_1^{-1} + m)^{-1}(1-\beta)} \right) \right]^{-1}}_{\mathcal{A}_4(\mathcal{B}_3, r, \beta, G_2, m, \xi_1)}. \quad (3.27)$$

Combining (3.22) and (3.27), we find that

$$\begin{aligned} \|\mathcal{F}_{1,\epsilon} - \mathcal{F}_{2,\epsilon}\|_{\mathcal{H}^j(\Omega)}^2 & \leq |\mathcal{A}_4(\mathcal{B}_3, r, \beta, G_2, m, \xi_1)|^2 \epsilon^2 \\ & \quad \times \sum_{\xi_n \leq \mathcal{C}_\epsilon} \xi_n^{2j+2} \left(\int_{\Omega} f(x) \varphi_n(x) dx \right)^2. \end{aligned} \quad (3.28)$$

The finite sum on the right above can be bounded as follows

$$\begin{aligned} \sum_{\xi_n \leq \mathcal{C}_\epsilon} \xi_n^{2j+2} \left(\int_{\Omega} f(x) \varphi_n(x) dx \right)^2 & \leq |\mathcal{C}_\epsilon|^2 \sum_{\xi_n \leq \mathcal{C}_\epsilon} \xi_n^{2j} \left(\int_{\Omega} f(x) \varphi_n(x) dx \right)^2 \\ & \leq |\mathcal{C}_\epsilon|^2 \|f\|_{\mathcal{H}^j(\Omega)}^2. \end{aligned} \quad (3.29)$$

Therefore, we follows from (3.28) that

$$\|\mathcal{F}_{1,\epsilon} - \mathcal{F}_{2,\epsilon}\|_{\mathcal{H}^j(\Omega)} \leq \mathcal{C}_\epsilon \epsilon \left| \mathcal{A}_4(\mathcal{B}_3, r, \beta, G_2, m, \xi_1) \right| \|f\|_{\mathcal{H}^j(\Omega)}. \quad (3.30)$$

Step 3. Estimate of $\|\mathcal{F}_\epsilon^1 - f_\epsilon\|_{\mathcal{H}^j(\Omega)}$. We derive that

$$\begin{aligned} & \mathcal{F}_{1,\epsilon}(x) - f_\epsilon(x) \\ &= \sum_{\xi_n \leq \mathcal{C}_\epsilon} \left[\mathcal{A}_{1,n}(m, \beta) \int_0^1 E_{\beta,\beta}(-\mathcal{A}_{2,n}(m, \beta)(1-s)^\beta)(1-s)^{\beta-1} g_\epsilon(s) ds \right]^{-1} \\ & \quad \times \left(\int_\Omega (\rho_\epsilon(x) - \rho(x)) \varphi_n(x) dx \right) \varphi_n(x). \end{aligned} \quad (3.31)$$

By taking the norm of both sides of the above expression in space $\mathcal{H}^j(\Omega)$, and using Parseval' s equality, we obtain that

$$\begin{aligned} & \|\mathcal{F}_{1,\epsilon} - f_\epsilon\|_{\mathcal{H}^j(\Omega)}^2 \\ &= \sum_{\xi_n \leq \mathcal{C}_\epsilon} \left[\mathcal{A}_{1,n}(m, \beta) \int_0^1 E_{\beta,\beta}(-\mathcal{A}_{2,n}(m, \beta)(1-s)^\beta)(1-s)^{\beta-1} g_\epsilon(s) ds \right]^{-2} \\ & \quad \times \xi_n^{2j} \left(\int_\Omega (\rho_\epsilon(x) - \rho(x)) \varphi_n(x) dx \right)^2. \end{aligned} \quad (3.32)$$

By looking back the inequality (3.26), we get

$$\begin{aligned} & \|\mathcal{F}_{1,\epsilon} - f_\epsilon\|_{\mathcal{H}^j(\Omega)}^2 \\ & \leq \left(\frac{m\mathcal{M}(\beta)}{\frac{\mathcal{M}(\beta)}{\xi_1\sigma_1} + (1-\beta)} \left[1 - E_{\beta,1} \left(- \frac{\beta(\xi_1^{-1} + m)^{-1}}{\mathcal{M}(\beta) + (\xi_1^{-1} + m)^{-1}(1-\beta)} \right) \right] \right)^{-1} \\ & \quad \times \sum_{\xi_n \leq \mathcal{C}_\epsilon} \xi_n^{2j+2} \left(\int_\Omega (\rho_\epsilon(x) - \rho(x)) \varphi_n(x) dx \right)^2. \end{aligned} \quad (3.33)$$

Form (3.33), one has

$$\begin{aligned} & \sum_{\xi_n \leq \mathcal{C}_\epsilon} \xi_n^{2j+2} \left(\int_\Omega (\rho_\epsilon(x) - \rho(x)) \varphi_n(x) dx \right)^2 \\ &= \sum_{\xi_n \leq \mathcal{C}_\epsilon} \xi_n^{2j+2+\frac{N}{r}-\frac{N}{2}} \xi_n^{\frac{Nr-2N}{2r}} \left(\int_\Omega (\rho_\epsilon(x) - \rho(x)) \varphi_n(x) dx \right)^2 \\ & \leq (\mathcal{C}_\epsilon)^{2j+2+\frac{N}{r}-\frac{N}{2}} \sum_{\xi_n \leq \mathcal{C}_\epsilon} \xi_n^{\frac{Nr-2N}{2r}} \left(\int_\Omega (\rho_\epsilon(x) - \rho(x)) \varphi_n(x) dx \right)^2 \\ &= (\mathcal{C}_\epsilon)^{2j+2+\frac{N}{r}-\frac{N}{2}} \|\rho_\epsilon - \rho\|_{\mathcal{H}^{\frac{Nr-2N}{4r}}(\Omega)}^2. \end{aligned} \quad (3.34)$$

Since $1 < r < 2$, with $L^r(\Omega) \hookrightarrow \mathcal{H}^{\frac{Nr-2N}{4r}}(\Omega)$. Therefore, we get

$$\|\rho_\epsilon - \rho\|_{\mathcal{H}^{\frac{Nr-2N}{4r}}(\Omega)} \leq \mathcal{C}_1(N, r) \|\rho_\epsilon - \rho\|_{L^r(\Omega)} \leq \mathcal{C}_1(N, r) \epsilon. \quad (3.35)$$

By summarizing all three evaluations (3.33), (3.34) and (3.35), we derive that

$$\|\mathcal{F}_{1,\epsilon} - f_\epsilon\|_{\mathcal{H}^j(\Omega)} \leq \mathcal{A}_5(\xi_1, \sigma_1, m, \beta) (\mathcal{C}_\epsilon)^{j+1+\frac{N}{2r}-\frac{N}{4}} \epsilon, \quad (3.36)$$

whereby

$$\begin{aligned} & \mathcal{A}_5(\xi_1, \sigma_1, m, \beta) \\ &= \left(\frac{m\mathcal{M}(\beta)}{\frac{\mathcal{M}(\beta)}{\xi_1\sigma_1} + (1-\beta)} \left[1 - E_{\beta,1} \left(- \frac{\beta(\xi_1^{-1} + m)^{-1}}{\mathcal{M}(\beta) + (\xi_1^{-1} + m)^{-1}(1-\beta)} \right) \right] \right)^{-1}. \end{aligned} \quad (3.37)$$

From (3.19) to (3.37), we can conclude that

$$\begin{aligned} \|f_\epsilon^{C_\epsilon} - f\|_{\mathcal{H}^j(\Omega)} &\leq |C_\epsilon|^{-k} \|f\|_{\mathcal{H}^{j+k}(\Omega)} + C_\epsilon \epsilon |\mathcal{A}_4(\mathcal{B}_3, r, \beta, G_2, m, \xi_1)| \|f\|_{\mathcal{H}^j(\Omega)} \\ &\quad + \mathcal{A}_5(\xi_1, \sigma_1, m, \beta) (C_\epsilon)^{j+1+\frac{N}{2r}-\frac{N}{4}} \epsilon. \end{aligned} \quad (3.38)$$

By using Lemma 2.7, since $0 < j < \frac{N}{4}$, we know that $\mathcal{H}^j(\Omega) \hookrightarrow L^{\frac{2N}{N-4j}}(\Omega)$, which yields to the desired result (3.38). \square

Acknowledgments. The authors would like to thank the anonymous referee for his/her comments that helped us improve this article.

REFERENCES

- [1] G.M. Bahaa, A. Hamiaz, *Optimality conditions for fractional differential inclusions with nonsingular MittagLeffler kernel*. Adv Differ Equ (2018), 257.
- [2] Paulo Mendes de Carvalho-Neto and Gabriela Planas, *Mild solutions to the time fractionalNavier-Stokes equations in R^N* , J. Differential Equations **259**, no. **7**, (2015), 29482980.
- [3] F. Mainardi, *Fractional calculus and waves in linear viscoelasticity*, Imperial CollegePress, London, 2010. An introduction to mathematical models.
- [4] M.M. Khader and K. M. Saad, *On the numerical evaluation for studying the fractional KdV,KdV-Burgers, and Burgers equations*, Eur. Phys. J. Plus **133** (2018), 113.
- [5] P. Chen and L. Y. Xiang, *Existence of mild solutions for fractional evolution equations with mixed monotone nonlocal conditions*, Z. Angew. Math. Phys. **65**, no. **4**, (2014), 711728.
- [6] Igor Podlubny, *Fractional differential equations*, Mathematics in Science and Engineering, Vol. **198**, Academic Press, Inc., San Diego, CA, 1999.
- [7] N.H. Luc, L.N. Huynh, D. Baleanu, & N.H. Can. Identifying the space source term problem for a generalization of the fractional diffusion equation with hyper-Bessel operator. Advances in Difference Equations, 2020, 1-23.
- [8] N.H. Can, O. Nikan, M.N. Rasoulizadeh, H. Jafari, & Y.S. Gasimov, Y. S., *Numerical computation of the time non-linear fractional generalized equal width model arising in shallow water channel*. Thermal Science, (2020), 49-58.
- [9] N. H. Tuan, A. T. Nguyen, and C. Yang, *Global well-posedness for fractional Sobolev-Galpern type equations*, DiscreteContin. Dyn. Syst. **42**, no. **6**, (2022), 26372665.
- [10] N. A. Tuan, N. H. Tuan, and C. Yang, *On Cauchy problem for fractional parabolic-elliptic Keller-Segel model*, Adv. Nonlinear Anal. **12** (2023), no. **1**, 97116.
- [11] N. H. Tuan, V. V. Au, and A. T. Nguyen, *Mild solutions to a time-fractional Cauchy problem with nonlocal nonlinearity inBesov spaces*, Arch. Math. **118**, no. **3**, (2022), 305314.
- [12] N. H. Tuan, M. Foondun, T. N. Thach, and R. Wang, *On backward problems for stochastic fractional reaction equations withstandard and fractional Brownian motion*, Bull. Sci. Math. **179**, no. **103-158**, (2022) 58 pp.
- [13] N. T. Bao, T. Caraballo, N. H. Tuan, and Y. Zhou, *Existence and regularity results for terminal value problem for nonlinear fractional wave equations*, Nonlinearity **34** (2021), 14481503.
- [14] N.D. Phuong, H.T.K. Van, N.H. Luc, L.D. Long. *Determine the unknown source term for a fractional order parabolic equation containing the Mittag-Leffler kernel*. Journal of Nonlinear and Convex Analysis, **23** (8), (2022), 1577-1600.
- [15] T. Caraballo, T.B. Ngoc, N.H. Tuan, & R. Wang. *On a nonlinear Volterra integrodifferential equation involving fractional derivative with Mittag-Leffler kernel*, Proceedings of the American Mathematical Society, **149**(08), (2021), 3317-3334.

- [16] N.H. Can, N.H. Luc, D. Baleanu, Y. Zhou Y. *Inverse source problem for time fractional diffusion equation with Mittag-Leffler kernel*. Advances in Difference Equations, **1**, (2020) 1-18.
- [17] N.H. Tuan, T. Caraballo. *On initial and terminal value problems for fractional nonclassical diffusion equations*. Proceedings of the American Mathematical Society, **149**(1),(2021), 143-161.
- [18] N.H. Tuan, T.N. Thach, N.H. Can, & D. O'Regan. *Regularization of a multidimensional diffusion equation with conformable time derivative and discrete data*. Mathematical Methods in the Applied Sciences, **44**(4), (2021), 2879-2891.
- [19] T.B. Ngoc, D. Baleanu, L.T.M. D., & N.H.Tuan, *Regularity results for fractional diffusion equations involving fractional derivative with MittagLeffler kernel*. Mathematical Methods in the Applied Sciences, **43**(12), (2020), 7208-7226.
- [20] F. S. A. Musalhi, S. A. S. Nasser and K. Erkinjon, *Initial and boundary value problems for fractional differential equations involving Atangana-Baleanu derivative*, SQU J. Sci. **23** (2018),137146
- [21] M.Q. Vinh, E. Nane, D. O'Regan, N.H. Tuan, *Terminal value problem for a nonlinear parabolic equation with Gaussian white noise*, Electronic Research Archive, Volume **30**, Issue **4**, (2020) 1374-1413.
- [22] N.H. Tuan, T. Caraballo, P.T.K. Van, V.V. Au. *On a terminal value problem for parabolic reactiondiffusion systems with nonlocal coupled diffusivity terms*, Communications in Non-linear Science and Numerical Simulation, (2022), 106248
- [23] N.H. Tuan, D. Lesnic, T.N. Thach, & T.B. Ngoc. *Regularization of the backward stochastic heat conduction problem*. Journal of Inverse and Ill-posed Problems, **30**(3), (2022), 351-362.
- [24] N.H. Tuan, N.M. Hai, T.N. Thach, & N.H. Can. *On stochastic elliptic equations driven by Wiener process with non-local condition*. Discrete and Continuous Dynamical Systems-S, (2022).

BUI DUC NAM

HO CHI MINH CITY UNIVERSITY OF INDUSTRY AND TRADE

E-mail address: nambd@hufi.edu.vn

LE XUAN DAI

VIETNAM NATIONAL UNIVERSITY, HO CHI MINH CITY, VIETNAM (VNU-HCM) AND DEPARTMENT OF APPLIED MATHEMATICS, FACULTY OF APPLIED SCIENCE, HO CHI MINH CITY UNIVERSITY OF TECHNOLOGY (HCMUT), 268, LY THUONG KIET, DISTRICT 10, WARD 14, HO CHI MINH CITY, VIETNAM

E-mail address: ytkadai@hcmut.edu.vn

LE DINH LONG

FACULTY OF MATHS, FPT UNIVERSITY HCM, SAIGON HI-TECH PARK, HO CHI MINH CITY, VIETNAM

E-mail address: longld13@fe.edu.vn

NGUYEN HOANG TUAN*

VIETNAM NATIONAL UNIVERSITY, HO CHI MINH CITY, VIETNAM (VNU-HCM) AND DEPARTMENT OF APPLIED MATHEMATICS, FACULTY OF APPLIED SCIENCE, HO CHI MINH CITY UNIVERSITY OF TECHNOLOGY (HCMUT), 268, LY THUONG KIET, DISTRICT 10, WARD 14, HO CHI MINH CITY, VIETNAM

E-mail address: Corresponding author : nhtuan.sdh231@hcmut.edu.vn